

W.T. Neale & Co., Ltd.,  
B.M. & S. Lamb  
16/18 Franklin Rd.  
Worthing, Sussex, BN132PQ,  
England

Harold Weiner  
Kaiserstr. 74  
3250 Hameln 1  
West Germany

World Insectivorous Plants  
2130 Meadowind Ln.  
Marietta, GA 30062

Wilson Yeo  
31 Seletar  
Singapore 2819

Inquire *Sarracenia, Dionaea, Darlingtonia seed*

Inquire *Aldrovanda, Byblis, Cephalotus, Dionaea, Drosophyllum, Drosera, Pinguicula, Sarracenia, Nepenthes*

\$.50 *Dionaea, Drosera, Drosophyllum, Nepenthes, Sarracenia, Pinguicula, Byblis liniflora, Utricularia*

Free *Malaysian Nepenthes species*

The co-editors and CPN do not endorse any of the above vendors. This is being provided only as a service to our subscribers. Information correct at time of receipt. Please contact sources directly for further information. Not responsible for omissions. Inquiries for inclusion in future lists should include a catalog/price list of CP available, cost of catalog and address. Send information to J.A. Mazrimas, 329 Helen Way, Livermore, CA 94550.

---

---

## Review of Recent Literature

# A NEW FAMILY OF CARNIVOROUS PLANTS

by J.A. Mazrimas

Givnish, T.J., E.L. Burkhardt,  
R.E. Happel, and J.D. Weintraub.

Carnivory in the Bromeliad *Brocchinia reducta*, with a cost/benefit model for the general restriction of carnivorous plants to sunny, moist, nutrient-poor habitats.

The American Naturalist 124, No. 4:  
479-497 1984

What really is a carnivorous plant? The authors of this paper propose a two part definition which is as follows:

The plant must be "able to absorb nutrients from dead animals juxtaposed to its surfaces, and thereby obtain some increment to fitness in terms of increased

growth, chance of survival, pollen production, or seed set.

Second, the plant must have some unequivocal adaptation or resource allocation whose primary result is the active attraction, capture, and/or digestion of prey."

For some time, observers of the tropical forest canopy noticed the cup-like structure (called a tank) of epiphytic bromeliads which provide the opportunity to store not only water but many forms of animal life which accidentally fall in and drown and eventually decay into simple nutrients. The question is whether these plants are benefiting by this passive capture which is the subject of this paper.

In the definition above, the first portion divides the plant kingdom into those plants that kill or immobilize an insect just for defensive purposes such as *Roridula gorgonias*. Here, the sticky tentacles trap the insect as a defensive measure. Eventually, the leaf falls off the plant and the insect decomposes in the soil at the base of the plant. The second part of the definition restricts this type of activity being recognized as carnivory since nematodes and earthworms eventually die and decompose into simple nutrients that plants utilize.

There are 16 recognized genera of carnivorous plants either on logical or historical reasons. Not all of them fulfill the above definition. Some lack the attractants for trapping prey (*Lentibulariaceae*) and others lack digestive glands and rely on bacteria for digestion (*Heliamphora* and *Darlingtonia*). Most Bromeliads fall into the category of saprophytes because they lack the means to attract, digest prey and the means to prevent insect escape.

The *Brocchinia* bromeliad grows on a sterile sandstone surface in the Gran Sabana in southeastern Venezuela. It grows on the tepuis, flat-topped highly acidic sandy soil areas, about 3,600 feet (120 Meters) and above the forest floor. Interestingly, it grows next to *Heliamphora*, *Drosera*, *Utricularia*, and *Genlisea* all of which are known carnivorous plants. It's a very common plant belonging to a genus with 18 known species and is found not only in Venezuela but also Guayana, but generally restricted to the Guyana Highlands.

What does the plant look like? The plant is about 12 inches (32 cm) in height and its yellowish-green leaves form a cylinder with the edges of the leaves overlapping tightly to retain water in the tank. The inside leaf surface is coated with a fine waxy powder which easily loosens when insects try to escape. The author demonstrates this with ants that failed to negotiate the vertical walls even though they could easily climb glass surfaces. The



*Brocchinia reducta* in habitat, Southeastern Venezuela.

Photos by T.J. Givnish, Harvard University.

fluid inside the tank is very acidic with a pH of 2.8-3.0. Microscopic examination of the plant showed no evidence for digestive glands.

The biomass within the tank was mostly ants (90%) and flies (5%) which is consistent with the fact that the tank fluid emits a sweet, nectarlike odor attractive to ants. This *Brocchinia* species is the only one of four others tested that had such a strong scent.

Trichomes, microscopic structures composed of specialized cells, are common in the Bromeliad family since they are capable of absorbing mineral nutrients essential to the well-being of epiphytic species. In *B. reducta*, these structures are very numerous at the inside base of the leaves and were shown to absorb amino acids in a radioactive form.

Protocarnivorous plants are those plants that have many of the preadaptations for attraction, capture, or digestion, but each of these traits may have other purposes with no clear allocation devoted solely to carnivory. Several species of Bromeliads fall into this category. The authors emphasize that the real crucial step for *B. reducta* is that it is capable of leaking a sweet smelling, volatile substance into its tank from glands in its leaf bases. They feel the key trait is attracting insects to the tank and this promoted the evolution of all the other functions that a true carnivorous plant must have.

This new representative of the carnivorous plant group is the least specialized of all the known carnivorous plants. It lacks digestive glands, specialized nectaries and has only a rudimentary waxy cuticle enclosing its water tank (lacks the bristles found in *Sarracenia*). Its traps are composed of single leaves formed into a rosette but retains the capacity to replace individual leaves as one dies which is not the case in *Heliamphora*.

This plant is an interesting case because it belongs to a genus in which not all the species are carnivorous and therefore affords a way of studying the various steps in evolution toward carnivory.

There are three factors, according

to the authors, that all carnivorous plants must have in their environment to survive:

- 1) A nutrient-poor soil
- 2) A sunny location
- 3) Seasonally moist soil

It's a fact that there are 60% less carnivorous epiphytic angiosperms than terrestrial types. *Nepenthes* (6 of 71 species) and *Utricularia* (12 out of 280 species) are epiphytic. This brings up another question: why are carnivorous plants carnivorous? There are 3 possibilities: First, the nutrients that are absorbed can elevate the rate and seasonal duration of photosynthesis. There is some evidence for this from controlled feeding experiments and careful measurements of the leaf mass. Secondly, carnivory increases flower and seed production. Thirdly, that mineral nutrition is partly replaced by organic nutrition. On this latter point, all available evidence points to the fact that mineral nutrition is the chief means by which carnivorous plants achieve growth. Possibility No. 2 could easily be inversely related to No. 1. The energy and nutrients from a few less leaves could easily be directed to producing more flowers or seed.

It seems rather elementary that in nutrient poor soils, carnivorous plants will do better in a sunny, moist habitat than in a shady or dry one provided there is no limit on nutrient availability. Epiphytes must also have a good source of moisture as *Nepenthes* vines do by contacting the moisture in the soil via roots. Epiphytic *Utricularia* grow in wet cloud forests where water stress is rare. However, bromeliads are frequently found on shady perches or if in a sunny location are usually subject to dessication. This may account for the low numbers of epiphytic carnivorous plants.

Another factor why epiphytes are selected against may be because of the benefits that these plants get from certain species of ants that protect the plant. The plant provides shelter and nectar and the plant receives in a passive manner a supply of nutrients into the various plant

recesses in the form of food wastes, dead nestmates, food storage, etc. When it is all added up, the benefits may outweigh the plant's ability or need to become carnivorous.

I am looking forward to the next article on this new member of the carnivorous plant family when the authors will go into detail on various aspects not covered in this report.

Kondo, K. and P.S. Lavarack. 1984. A cytotaxonomic study of some Australian species of *Drosera* L. (Droseraceae). Bot. J. Linnean Soc. 88:317-333.

Fifteen species are presented along

with eleven new chromosome counts. *Droseras* in Australian present a large aneuploid series. The lowest chromosome number is *D. paleacea* with  $2n=10$ , giving a new basic number  $x=5$  for the genus. Several rearrangements within subsections of the genus are proposed.

DES

Taylor, Peter. 1984. Two new bladderworts from South India. Proc. Indian Acad. Sci. (Plant Sci.) 93:99-103.

Two small annual terrestrial Utricularias are herein described for the first time for South India: *U. ceceili* and *U. lazulaina* with complete botanical descriptions and line drawings. DES

## WANT ADS

When submitting Want Ads, please be sure to print clearly for best results and to eliminate mistakes. Please circle the correct letter before each item (Want, Trade, Sell or Buy). Want ads are limited to carnivorous plants, terrariums, greenhouses and moss. There is a charge of ten cents per item, with no limit to the number of items you may submit per issue.

Send coin or check to:  
Arboretum, Want Ads  
California State University  
Fullerton, CA 92634

William DiLapi (Thompson Rd., Oswego, NY 13126) (WTB) any Pygmy *Drosera* plants or gemmae. (WTB) any terrestrial *Utricularia*. Send lists available.

Austin Gavin (7913 Behtelen Woods Lane, Springfield, VA 22153) (WTB) cuttings of *N. lowii* or seeds. (T) *N. rafflesiana*, *N. ventricosa* or *Cephalotus*.

Peter Harle (418 Point Mary, Dunkirk, MD 20754) (WB) *S. minor* × *leucophylla* (plants); *D. adelae*; *Heliophora* plant; small, dome-type terrarium.

Matt Hochberg (5500 Fieldston Rd., Bronx, NY 10471) (B)(W) 1. Any *Sarracenia* hybrids, (B)(W) 2. Any *Drosera* (except *D. capillaris*), (B)(W) 3. Any other carnivorous plants. (Send me your price list of plants/cuttings/seedlings available.)

Lee's Botanical Gardens (P.O. Box 7026, Ocala, FL 32672) (TS) 100 varieties of CP, (WT) Will

exchange generously for *N. × liveyi*, *Genlisea*, terrestrial and epiphytic orchids, staghorn ferns, free list — inquire.

Claus Thiede (Goslarsche Str. 70, D-3300 Braunschweig, W-Germany) has *N. albo-marginata* (seed/plants), *N. pectinata*, *N. tobaica* (seed), *N. hookeriana* (F1), *N. treubiana* and other *Nepenthes* plants/seed from Borneo, Sumatra, Malaysia for other comparable uncommon *Nepenthes* plants (no hybrids) more for trade than to sell.

Harald Weiner (Kaiserstr. 74, 3250 Hameln 1, West Germany) Trade: *Pinguicula crystallina*, *P. hirtiflora*, *P. zecheri*, *P. ehlersae*, *P. esseriana*, *P. vallisneriaefolia*, *P. filifolia*, *Drosera arenicola*, *D. roraimae*, *D. Collinsiae*, *D. platypoda*, *Nepenthes madagascariensis*, *N. pervillei*, *N. truncata* and various other CP. Want: *Pinguicula macrocerus*, *P. m. nortensis*, *P. ramosa*, *Drosera ramentacea*, *D. affinis*, *D. alba*, *D. felix*, *D. graminifolia*, *Sarracenia flava* all red type, *S. alabamensis*, *Nepenthes clipeata*, *N. singalana*, *N. bongso*, *N. pectinata* and others.

Andreas Wistuba (Mudauer Ring 227, D-6800 Mannheim 52, West Germany) (TS) Many interesting CP including *Drosera arenicola*, *D. caledonica*, *D. hamiltonii*, *D. intermedia* 'Roraima', *D. prolifera*, *D. roraimae*, New South African *Drosera* species, *Pinguicula ehlersae*, *P. esseriana*, *Sarracenia* seedlings, *Utricularia livida*, (WTB) *Drosera acaulis*, *D. banksii*, *D. cyemensis*, *D. petiolaris* (different types), *D. ramentacea*, any other rare CP, South African, South American and Australian CP. (WTB) Seeds of rare *Sarracenia* forms and hybrids, *Biovularia*, *Genlisea*, *Heliophora*, Mexican *Pinguicula* species, (W) Contacts to CP collectors all over the world.



# CARNIVOROUS PLANT NEWSLETTER

Official Journal of the  
International Carnivorous  
Plant Society



Volume 14, Number 1  
March 1985

## COVER PHOTO

*Brocchinia reducta*, a bromeliad newly described as carnivorous. Photo by T.J. Givnish, Harvard University. See article, page 24.

The co-editors of CPN would like everyone to pay particular attention to the following policies regarding your dues to the ICPS.

All correspondence regarding dues, address changes and missing issues should be sent to Mrs. Pat Hansen, 3321 Hamell Rd., Fullerton, CA 92635. DO NOT SEND TO THE CO-EDITORS. Checks for subscriptions and reprints should be made payable to CSUF FOUNDATION-ICPS.

All material for publication, comments and general correspondence about your plants, field trips or special noteworthy events relating to CP should be directed to one of the co-editors. We are interested in all news related to carnivorous plants and rely on the membership to supply us with this information so that we can share it with others.

Views expressed in this publication are those of the authors, not necessarily the editorial staff.

Copy deadline for the September issue is July 1, 1985.

### CO-EDITORS:

D.E. Schnell, Rt. 1, Box 145C, Pulaski, VA 24301

J.A. Mazrimas, 329 Helen Way, Livermore, CA 94550

T.L. Mellichamp, Dept. of Biology, UNCC, Charlotte, NC 28223

Leo Song, Dept. of Biology, California State University, Fullerton, CA 92634

Seed Bank: Patrick Dwyer, St. Michael's Episcopal Church, 49 Killean Park, Albany, NY 12205, USA.

BUSINESS MANAGER: Mrs. Pat Hansen, c/o The Fullerton Arboretum

PUBLISHER: The International Carnivorous Plant Society by the Fullerton Arboretum, California State University, Fullerton, CA 92634. Published quarterly with one volume annually. Printer: Kandid Litho, 129 Agostino Rd., San Gabriel, CA 91776. Circulation: 322 (68 new, 254 renewal). Dues: \$10.00 annually, \$15.00 foreign. Reprints available by volume only. ©1985 Carnivorous Plant Newsletter. All rights reserved.

# ***CARNIVOROUS PLANT NEWSLETTER***

VOLUME 14, Number 1

MARCH 1985

