

HELIAMPHORA CHIMANTENSIS, A NEW SPECIES OF *HELIAMPHORA* (SARRACENIACEAE) FROM THE 'MACIZO DE CHIMANTA' IN THE SOUTH OF VENEZUELA

ANDREAS WISTUBA • Mudauer Ring 227 • 68259 Mannheim • Germany
THOMAS CAROW • Am Mustergarten 1 • 97702 MÜNNERSTADT • Germany
PETER HARBARTH • Frankenweg 18 • 69221 Dossenheim • Germany

Keywords: new taxa: *Heliamphora chimantensis*, Venezuela.

Received: 8 February 2002

Introduction

During our January 2001 expedition to the tepuis of Venezuela (Wistuba *et al.*, 2001), we also explored parts of the 'Macizo de Chimanta', the Chimanta Massif in the southwest part of the Gran Sabana. This huge massif covers a total area of 1470 km² and is actually a cluster of tepuis including the central Chimanta Tepui itself. Their peaks range in altitude from 1700 m (at the central part of the massif) to 2698 m (on Eruoda Tepui). The ten tepuis that reach 2000 m above sea level cover an area of some 700 km² in total (Huber, 1992). The size and the diverse altitudes of the Chimanta Massif support numerous habitats including rivers, green valleys and forests as well as rocky plateaus and moist savannas. Starting in the 1950s, various expeditions explored the unique flora and fauna of this area. Many endemic plants and animals were discovered during these expeditions and certainly many more await discovery.

During this exploration we found a previously undescribed species of *Heliamphora* most notable for its pitcher shape and the spoon shaped lids. This species seems to be more closely related to *Heliamphora tatei* than to any of the other species described so far from the Gran Sabana.

Heliamphora chimantensis Wistuba, Carow & HarbARTH *spec. nov.*

Caudex ramosus; foliis caulescentibus vel rosulis; amphoriis in parte inferiore infundibuliformibus, in parte media subventricosis, et in parte superiore cylindricis ad leviter infundibuliformibus, longis 20-30 cm, latis (in parte superiore) 3.5 - 5 cm.

Inflorescentiis 3-5-floris, racemosis, ad 65 cm longis; flores nutantes; pedicellis 5-13 cm longis; petalis 4 lanceolatis, albidis vel pallide-roseis, 4-5.5 cm longis, 1.2-1.7 cm latis; staminibus ca. 20, 1-serialibus, filamentis 6 mm longis; antheris oblongo-lanceolatis, ca. 5 mm longis; ovario valde tomentoso; stylo glabro; stigmatibus 3 lobato; seminibus fuscis, oblongis, ca. 3 mm longis, testa conspicue membranaceo-alata.

Rhizomes branching. Plants forming dense and often huge patches up to several meters across.

Pitchers infundibulate in the lower half, slightly ventricose in the middle and cylindrical to slightly infundibulate in the upper third (see Figure 1); pitchers 20 to 35 cm long, 3.5-5 cm wide in the upper part; upper part of the pitchers completely glabrous on the inner side; pitchers entirely green with deep red lids. Lid 1-2 cm wide and 2-2.5 cm long, spoon-shaped, upright, ending with a sharp tip; the two lobes of the lid compressed from the sides near the tip, often touching each other at the front, forming a quasi-helmet; lobes are expanded in the lower part of the lid and narrowed sharply near the base; the inner side of the lid with prominent irregularly shaped patches of glands, up to 5 mm across. Inflorescence 60 to 65 cm long, 3-5 flowers, peduncle slightly pubescent,

pedicels 5-13 cm long. The lowest peduncle is the longest and bears a bract that frequently is transformed into a rudimentary pitcher. Tepals lanceolate, 4.5-5.5 cm long, 1.2-1.7 cm wide, white to whitish-pink; ca. 20 stamens in 1 series, filaments 6 mm long, anthers oblong lanceolate, 5 mm long, 1 mm wide; ovary 3 celled, pubescent, style glabrous. Seed approximately 2 mm long, compressed, ovate, irregularly winged.

Specimens examined

Heliamphora chimantensis: Macizo de Chimanta, Section centro-oriental, 1921 m a.s.l., N 05°16.672'; W 062°11.438' Wistuba, Carow & Harbarth No. Chim 10.01.01/1, holotype, flowering plant (VEN)

Heliamphora chimantensis: Macizo de Chimanta, Section centro-oriental, 1921 m a.s.l., N 05°16.672'; W 062°11.438' Wistuba, Carow & Harbarth No. Chim 10.01.01/2, isotype, flowering plant (VEN)

The two herbarium specimens nicely exhibited the characteristics typical of the new species. The comments in this paper are based both upon these two specimens as well as our field observations of many other plants in situ.



Figure 1: *Heliamphora chimantensis* pitcher. Photograph by Andreas Wistuba.



Figure 2: Cluster of *Heliamphora chimantensis* pitchers. Photograph by Andreas Wistuba.



Figure 3: *Heliamphora chimantensis* pitcher appendage nectar patches (left inset) and insect visitation; ant (left), wasp (right). Photographs by Thomas Carow.

Heliamphora chimantensis sometimes grows together with *H. minor*, the only other species of *Heliamphora* recorded from Chimanta. As a consequence, hybrids frequently could be found. However, *H. chimantensis* seems to prefer valleys growing at around 2000 m while *H. minor* also has been found in higher altitudes and usually prefers more open habitats.

The *H. minor* plants on Chimanta are notable for the long and prominent bristles inside the pitchers. This variant form of *H. minor* is actually fairly widespread and has been found on many tepuis of the Chimanta Massif as well as on Aprada Tepui, but not Auyan Tepui. Meanwhile, the typical form of *H. minor* is only known from Auyan Tepui. The differences between these two forms of *H. minor* may merit further taxonomic study.

Ecology

Heliamphora chimantensis plants were found growing exposed, in short vegetation such as grasses, bromeliads, *Xyris* (Xyridaceae) or *Stegolepis* (Rapateaceae). These plants do not grow taller than the *H. chimantensis*. In fact, we never found plants growing in shaded locations. Often they grow near rivers in the valleys of Chimanta. In all cases we found *Heliamphora chimantensis* growing together with *Stegolepis ligulata*, and also often associated with *Adenanthe bicarpellata* (Ochnaceae) and various species of *Brocchinia* (Bromeliaceae).

In comparison to the species found on the plateaus of the various tepuis, where the surface usually is much more rocky and sandy and plants often can grow only on 'islands' of debris, highly limited in space, the moist savannah-like habitat *H. chimantensis* prefers allows the formation of huge clumps. We have visited many tepuis on this and other expeditions, and had never before seen clumped *Heliamphora* colonies of comparable in size to the ones typical of *H. chimantensis*. Vegetative reproduction seems to play an important role as the seedling activity we observed was very low.

The lids of *H. chimantensis* bear strange, huge patches (up to 5 mm across) of glands on their inner surface (Figure 3). They should be studied in more detail as they seem to be involved in the attraction of prey. We observed ants and wasps being attracted by secretions of these glands but unfortunately our limited time on Chimanta did not permit a detailed study. As we have observed with other species of *Heliamphora*, the pitchers usually do not contain many captured insects.

Related species

The discovery of *H. chimantensis* on Chimanta and its characteristics came as quite a surprise, as they clearly indicate that it is much more closely related to the southern *H. tatei*, than to any of the northern species known to be growing in the Gran Sabana. The flowers of all other species known from the Gran Sabana have 10-15 anthers, while *Heliamphora tatei* var. *tatei* and *H. tatei* var. *neblinae* from the Amazon, and *Heliamphora chimantensis* have about 20; however, while the anthers of *H. tatei* and *H. tatei* var. *neblinae* are 7-9 mm long, those of *H. chimantensis* just reach 5 mm in length (Maguire, 1978; Steyermark, 1984).

The nectar spoons of the other species from the Gran Sabana are shaped more or less concave to form helmet-like structures under which nectar can accumulate, protected from the frequent rainfalls. In *H. tatei*, as well as in *H. chimantensis*, the lids are more upright. However, *H. chimantensis* protects its nectar production by the two lobes of the lid which are bent forwards and are compressed from the sides to form a roof reminiscent of the helmets of the other north-eastern species. From the ones of *H. tatei* the lids of *H. chimantensis* differ by the spoon-shape and the sharp contraction at the base. In *H. tatei* the lids are only very slightly contracted at the base and rather rectangular in shape.

We observed numerous hybrids between *Heliamphora chimantensis* and the *Heliamphora minor* on Chimanta. These hybrids can be easily distinguished from pure *Heliamphora chimantensis* by the shorter pitchers, the red veins and the coarse bristles inside the pitchers. Some of the hybrids' pitchers showed deep red coloration, similar to the form of *H. minor* growing on Chimanta. The helmet shaped lids of these show a strong influence of *Heliamphora minor*. Apparently the hybrids are much more vigorous than either of the parent plants; we found patches of single clones measuring more than 5 meters across (see Front Cover).

Etymology

This name was chosen as *Heliamphora chimantensis* because it is the only species known to be restricted to Chimanta Tepui.

Discussion

The formation of huge clumps and the low seedling activity seem to be the consequence of the savanna-like habitat. In general, the seeds of *Heliamphora* clearly show adaptations to dispersal by water. This works excellently on a "typical" tepui where little streams and shallow ponds carry the seeds to new small patches of debris, making *Heliamphora* a kind of pioneer plant. However, the environment in the valleys on Chimanta where we found *Heliamphora chimantensis* seems to be far more static, with a well developed, continuous soil layer. The low vegetation might be too dense for seedlings to germinate and grow. Those few seedlings which do survive to maturity can develop into patches of huge size.

Although the large sizes of the clumps that *Heliamphora chimantensis* makes are remarkable, they are not completely unprecedented. *Heliamphora* can multiply vegetatively by division, but in most species the resulting clumps still remain small due to limitations of the detritus patches they grow in. Even so, we have observed similar, large-clump growth patterns in *Heliamphora hispida* (Nerz & Wistuba, 2000), *Heliamphora heterodoxa* (in the Gran Sabana) and *Heliamphora nutans* growing on the sandy foothill area of Tramen Tepui.

As we already discussed in our publication on *Heliamphora folliculata* (Wistuba *et al.*, 2001) the pitcher-lids or so called nectar-spoons of *Heliamphora* are highly elaborate structures for the attraction of prey. Apparently various strategies have been developed by the different species of *Heliamphora* all using the lids in altered ways. *Heliamphora chimantensis* with its gland-patches shows yet another way to use the lid-structure for the attraction of insects.

Being morphologically fairly constant organs, the lids offer superb, yet previously under-utilized, characters of taxonomic relevance (cf. Nerz & Wistuba, 2000; Wistuba *et al.*, 2001).

Literature

- Huber, O. 1992, El macizo del Chimantá, Oscar Todtmann Editores.
 Maguire, B. 1978, Sarraceniaceae (*Heliamphora*), in: The Botany of the Guyana Highland Part-X, Memoirs of the New York Botanical Garden, 29: 36-61.
 Nerz, J. and Wistuba, A. 2000, *Heliamphora hispida* (Sarraceniaceae), a New Species from Cerro Neblina, Brazil-Venezuela. Carniv. Pl. Newslett. 29: 37-41.
 Steyermark, J.A. 1984, Venezuelan Guyana, Annals of the Missouri Botanical Garden, 71: 302-312.
 Wistuba, A., Harbarth, P. and Carow, T. 2001, *Heliamphora folliculata*, a New Species of *Heliamphora* (Sarraceniaceae) from the 'Los Testigos' Table Mountains in the South of Venezuela. Carniv. Pl. Newslett. 30:120-125.

CARNIVOROUS PLANT NEWSLETTER



Journal of the International
Carnivorous Plant Society
www.carnivorousplants.org

Volume 31, Number 3
September 2002



Front Cover: *Heliamphora chimantensis* × *Heliamphora minor* plants. Photograph by Andreas Wistuba. Article on page 78.

Back Cover: *Pinguicula moranensis* 'Libelulita' flowers, photo by Barry A. Rice. Article on page 83.

Carnivorous Plant Newsletter is dedicated to spreading knowledge and news related to carnivorous plants. Reader contributions are essential for this mission to be successful. Do not hesitate to contact the editors with information about your plants, conservation projects, field trips, or noteworthy events. Contributors should review the "Instructions to Authors" printed in the March issue of each year. Advertisers should contact the editors. Views expressed in this publication are those of the authors, not the editorial staff.

All correspondence regarding dues, address changes and missing issues should be sent to the Membership Coordinator at the ICPS. Do not send such correspondence to the editors. Checks for subscriptions and back issues should be made to the ICPS in US funds. Dues for 2002 are \$25.

ICPS, Inc.
PMB 330
3310 East Yorba Linda Blvd.
Fullerton, CA 92831-1709, USA
icps@carnivorousplants.org

President
Vice President
Secretary
Seed Bank

David Gray, email: david@carnivorousplants.org
Carl Mazur, email: carl@carnivorousplants.org
Cindy Slezak, email: cindy@carnivorousplants.org
John Brittnacher, email: john@carnivorousplants.org, seedbank listed in this issue.

Editors:

Barry A. Rice, P.O. Box 72741, Davis, CA 95617, USA, email: barry@carnivorousplants.org
Jan Schlauer, Zwischenstr. 11, D-60594 Frankfurt, Germany, email: jan@carnivorousplants.org
Page Layout: Steve Baker, email: steve@carnivorousplants.org

Date of effective publication of the June 2002 issue of Carnivorous Plant Newsletter: 6 June 2002.

The ICPS is the International Cultivar Registration Authority (ICRA) for cultivated carnivorous plants according to The International Code For The Nomenclature of Cultivated Plants. Send relevant correspondence to the ICPS, Inc.

PUBLISHER: ICPS, Inc., Fullerton, California. Published quarterly with one volume annually. Desktop Publishing: Steve Baker, 5612 Creek Point Drive, Hickory, NC 28601. Printer: Kandid Litho. Logo and masthead art: Paul Milauskas. Dues: \$25.00 annually. Reprints available by volume only © 2002 Carnivorous Plant Newsletter. All rights reserved. ISSN #0190-9215.

CARNIVOROUS PLANT NEWSLETTER

Journal of the International Carnivorous Plant Society

Volume 31, No. 3

September 2002

