

NEPENTHES ALBA AND *NEPENTHES GRACILLIMA*

STEWART MCPHERSON • Dorset, Poole • UK • stewart@redfermnaturalhistory.com

Keywords: *Nepenthes alba*, *Nepenthes gracillima*, Mount Tahan, Malaysia, taxonomy, Ridley.

One century ago, the English botanist Henry Ridley traveled to Malaysia and climbed Mount Tahan in (what is now) Pahang. During his climb, Ridley discovered two species of pitcher plants (*Nepenthes*). He described the first one under the name *Nepenthes gracillima* in 1908, and recorded that it produces long, slender pitchers that are predominantly black mottled with green blotches (Ridley 1908). Sixteen years later, Ridley (1924) described the second species which he had encountered on Mount Tahan, and he named that plant *N. alba* for it produces smaller, predominantly white pitchers.

During the one hundred years since Ridley's work, the classification of these two plants has been clouded with errors and confusion. Various botanists have considered the two plants the same species; others mistakenly identified either or both under various incorrect names, including *N. ramispina*, *N. bongso*, and *N. singalana*. Perhaps, as a result of this confusion, *Nepenthes alba* has remained absent from all recent listings of *Nepenthes* species, most notably Clarke (2001) and Rice (2006).

In 2008, I travelled to Mount Tahan to attempt to understand the taxonomy of the *Nepenthes* of that mountain. I observed populations of two distinct types of *Nepenthes* on the upper slopes of Mount Tahan, both of which were consistent with Ridley's original descriptions of *N. gracillima* and *N. alba*. The pitchers of these plants differ in shape, size, and colouration, and they predominantly occur at different altitudes and in distinct habitats. As far as I observed, they do not hybridize or introgress and are easily distinguished from one another. On the basis of my observations, I concluded that Ridley's original classification of these two plants as distinct species is correct, and *N. alba* and *N. gracillima* are indeed different species.

Nepenthes alba is named after the Latin *albus* (white). The name refers to the colouration of the upper pitchers of this plant (see Figure 1 and Front Cover). *Nepenthes alba* is widespread across the upper slopes of Mount Tahan, in Taman Negara (Malaysia's largest national park) and it occurs with increasing abundance from 1600 m (5250 ft) altitude to the summit of the mountain, which stands at 2187 m (7175 ft). The population of *N. alba* on Mount Tahan consists of many tens of thousands of plants which display consistent morphology, size, and general colouration (although the upper pitchers vary in colour). The wider distribution of *N. alba* beyond Mount Tahan remains unclear but herbarium specimens from Mount Tapis may represent



Figure 1: The pitchers of *Nepenthes alba* may be pure white.

this species too.

At lower elevations, *N. alba* grows amidst stunted, upper montane forest and scrub, mainly on ridge tops and the upper slopes of the mountain. It produces a branched stem up to 5 m (16 ft) long and scrambles and climbs amidst surrounding vegetation, and mostly grows in strong or direct sunlight. At higher elevations (especially on the upper slopes of Mount Tahan), it grows amidst stunted scrub or in exposed areas, devoid of vegetation, and often forms a compact rosette or short scrambling stem that trails across the ground. On the summit of the mountain, large populations of *N. alba* grow in waterlogged, marshy areas, amidst short marsh grasses and orchids. In such exposed conditions, the *N. alba* plants are greatly dwarfed and flowering specimens just 25 cm (10 in) tall are often observed (see Figure 5).

The lower pitchers of *N. alba* – note the typical purple colouration (see Figure 2). The lower pitchers are up to 12 cm (4.7 in) long and 4.5 cm (1.8 in) wide (although usually are much smaller than these maximum measurements). The bottom half of the pitcher is ovate and slightly swollen. Above this part, the width of the pitcher narrows (usually with a distinct hip) and the rest of the trap is cylindrical or very slightly infundibular towards the pitcher opening. Wings up to 7 mm (0.28 in) wide, fringed with filaments up to 10 mm (0.39 in) long run down the front of the lower pitchers. The peristome is up to 8 mm (0.31 in), and is lined with ribs up to 1 mm (0.04 in) high, spaced up to 1 mm (0.04 in) apart. The peristome is cylindrical and narrow at the front of the pitcher opening, and expanded and flattened at the back (especially below the lid). The peristome ribs are elongated along the inner edge of the peristome and form spikes up to 2 mm (0.08) long, which extend into the pitcher opening. The lid is sub-orbicular or elliptic, up to 4 cm (1.6 in) wide, 3.8 cm (1.5 in) long and lacks an appendage. The spur is generally unbranched and up to 13 mm (0.51 in) long.

The colouration of the lower pitchers is very consistent. The exterior surface of the pitcher is dark purplish brown, often mottled with faint dark purple blotches. The interior of the trap is uniformly light purple, sometimes with faint purple blotches. The lid (both sides) is light purple or reddish. The peristome is dark purple or black.

The upper pitchers are up to 13 cm (5 in) long and 4 cm (1.6 in) wide (although usually are much smaller than these maximum measurements). The bottom half of the pitcher is narrowly infundibular or ovate. Above this part, the width of the pitcher narrows (usually with a distinct hip) and is cylindrical towards the pitcher opening. Wings are reduced to narrow ridges which run down the corners of the flattened front face of the trap. The peristome is up to 7 mm (0.28 in) wide and is slightly expanded at the sides and back of the pitcher opening, or not at all. The peristome is lined with fine ribs up to 0.5 mm (0.02 in) high, spaced up to 1 mm (0.04 in) apart (although often the peristome ribs may be hardly discernable at all). All other parts are consistent with the lower pitchers.

The exterior and interior of the upper pitchers is predominantly white in colour, although the bottom half of the trap may suffuse yellowish green towards the base. Red or pink blotches and flecks are usually present either on the interior or exterior of the upper pitchers (or on both surfaces) although the extent and density of this colouration is extremely variable (see below). The



Figure 2: The lower pitchers of *N. alba* – note the typical purple colouration. Photo by Stewart McPherson

vestigial wing ridges which run down the front of the pitcher may be dark purple or may have no colouration at all in plants which produce pure white pitchers. The peristome is usually pure white, sometimes striped with bands of red, or suffused pinkish. The lid may be pure white, or mottled with red or pink blotches, or suffused reddish.

The colouration of the upper pitchers of this species is extremely variable. From my observations, I would estimate that approximately 90% of *N. alba* plants produce pitchers that are predominantly white with some red or pink blotches on the exterior or interior of the trap. About 9% of the total wild population on Mount Tahan produces upper pitchers which are consistently pure white. The remaining 1% produce upper pitchers that are very heavily lined with red blotches to the extent that they appear predominantly pure red, pink or orange in colour.

Nepenthes gracillima derives its name from the Latin *gracillimus* (slenderest or slimmest), which refers to the shape of the upper pitchers (see Figure 3). *Nepenthes gracillima* is currently known only from the upper slopes of Mount Tahan, and it seems to be much rarer than *N. alba*. All known populations (as well as my observations) are recorded amidst upper montane scrub on humid ridge tops between 1600-1700 m (5250-5577 ft) altitude. *Nepenthes gracillima* is not widespread or populous on Mount Tahan, and the populations I observed consisted of just a few dozen plants (but these are consistent and uniform).

Nepenthes gracillima readily forms a branched stem up to 3 m (10 ft) long, and scrambles and climbs amidst surrounding vegetation, particularly in areas where it may grow in strong or direct sunlight. In contrast to *N. alba*, *N. gracillima* does not grow on the ground in open, exposed habitat or grow in exposed areas and form stunted, compact foliage.

The lower pitchers are up to 22 cm (8.7 in) long and 7 cm (2.8 in) wide (see Figure 4). The bottom half of the pitcher is ovate and broad. Above this part, the trap narrows slightly (or hardly at all) with a faint hip (or no hip at all) and becomes cylindrical towards the pitcher opening. Wings up to 9 mm (0.35 in) wide fringed with narrow filaments up to 1 cm (0.39 in) long run down the front of the pitcher. The peristome is glossy, up to 1.8 cm (0.71 in) wide, and is narrow at the front of the pitcher opening, but expanded at the sides and back, especially below the lid. A gap a few millimeters wide may be present in the back of the peristome, immediately below the lid. The peristome with conspicuous ribs up to 1.5 mm (0.06 in) high, spaced up to 1.5 mm (0.06 in) apart. The inner margin of the peristome extends into the pitcher opening for several millimeters. The peristome ribs are elongated along the inner edge of the peristome and form spikes up to 2 mm (0.08 in) long, which extend into the pitcher opening. The lid is elliptic or ovate, up to 6 cm (2.4 in) long and 5 cm (2.0 in) wide and lacks an appendage. The spur is usually unbranched and up to 2 cm (0.8 in) long. The exterior of the lower pitcher is predominantly yellowish green, with variable, dark red or purple blotches. The peristome is green or reddish,



Figure 3: An upper pitcher of a plant growing on Mount Tahan that matches Ridley's description of *N. gracillima* – note the morphological differences with the upper pitchers of *N. alba*. Photo by Stewart McPherson

often striped with bands of dark red or purple. The interior of the pitcher and lower surface of the lid are yellowish green or light yellow. The upper surface of the lid is olive green, with variable dark purple or black flecks. In all populations of *N. gracillima* which I observed, the colouration of the lower pitchers was consistent (as described above).

The upper pitchers are up to 26 cm (10 in) long and 5 cm (2 in) wide (many times larger than the typical size of the upper pitchers of *N. alba*). The traps are wholly infundibular, although extremely narrowly so, and greatly elongated. Sometimes, the bottom third of the pitcher may be slightly swollen, with a very faint hip above. Wings are reduced to narrow ridges which run down the corners of the abruptly flattened, angular front face of the trap. The peristome is up to 12 mm (0.47 in) wide and is slightly expanded at the sides and back of the pitcher opening, or not at all. The peristome is lined with fine ribs up to 0.5 mm (0.02 in) high, spaced up to 1 mm (0.04 in) apart (although often the peristome ribs may be hardly discernable at all). All other parts are consistent with the lower pitchers.

The exterior of the upper pitcher is predominantly black, mottled with variable bright green blotches (which are elongated towards the base of the trap). This colouration is also present on the upper surface of the lid, however the lower surface of the lid and interior of the pitcher are light yellowish green. The peristome is yellow, striped with narrow bands of black and purple.

Recent observations of these two spectacular and very beautiful pitcher plant species on Mount Tahan raises many questions as to whether both also occur widely across the many peaks of central Peninsula Malaysia, and beyond. However, since both *N. alba* and *N. gracillima* occur within responsibly and strictly managed protected areas, both are not currently threatened with extinction. Certainly more extensive studies of these two taxa are required to understand their association and taxonomy.

An extensive account of the discovery, taxonomy, and morphology of *Nepenthes alba* and *N. gracillima* is presented in my new, two volume, 1399-page work *Pitcher Plants of the Old World*, which examines all species of *Nepenthes* known worldwide (see www.red-fernnaturalhistory.com for more information).



Figure 4: A lower pitcher of *N. gracillima* – note the profoundly different morphology (and size) in comparison to *N. alba*. Photo by Stewart McPherson




Figure 5: A stunted, flowering, *N. alba* plant growing on the bleak, windswept summit of Mount Tahan. Photo by Stewart McPherson.

References:

- Clarke, C.M. 2001. *Nepenthes* of Sumatra and Peninsular Malaysia. Natural History Publications (Borneo), Kota Kinabalu, Malaysia.
- Rice, B. 2006. *Growing Carnivorous Plants*. Timber Press, Portland, Oregon, USA.
- Ridley, H.M. 1908. On a collection of plants made by H. C. Robinson and L. Wray from Gunong Tahan, Pahang. *The Journal of Linnean Society, Botany*, 38: 320.
- Ridley, H.M. 1924. *Nepenthaceae*. *The Flora of the Malay Peninsula* 3: 21–25.

When in Northern California Visit



California Carnivores

Specializing in insect-eating and other exotic plants

- ★ Open all year, call ahead in winter
- ★ Commercially cultivated
- ★ Over 500 varieties on display
- ★ On site & domestic mail order
- ★ We ship potted plants

2833 Old Gravenstein Hwy South, Sebastopol, CA 95472
(707) 824-0433 Price List – SASE
Visit us on the Internet at <http://californiacarnivores.com>



petflytrap

THE ORIGINAL VENUS FLYTRAP

- Mail order and Internet U.S. sales
- *Dionaea*, *Sarracenia*, *Nepenthes*, *Drosera*, *Pinguicula*, books, supplies, and **much** more
- We stock **multiple sizes** of most plants
- Terrarium kits with **live plants** (no seeds)
- Rare nursery-grown, imported lowland/intermediate *Nepenthes*
- Most plants shipped potted and **ready to grow!**

www.petflytrap.com
orders@petflytrap.com
281-433-3286

CARNIVOROUS PLANT NEWSLETTER

Journal of the International Carnivorous Plant Society

Volume 38, No. 4

December 2009



CARNIVOROUS PLANT NEWSLETTER



Journal of the International
Carnivorous Plant Society
www.carnivorousplants.org

Volume 38, Number 4
December 2009



Front Cover: The spectacular pitchers of *Nepenthes alba* growing on the upper slopes of Mount Tahan. Photo by Stewart McPherson. Article on page 102.

Back Cover: The opening of a lower/intermediate pitcher of *Nepenthes attenboroughii* growing on the summit of Mount Victoria, Palawan. Photo by Stewart McPherson. Article on page 100.

Carnivorous Plant Newsletter is dedicated to spreading knowledge and news related to carnivorous plants. Reader contributions are essential for this mission to be successful. Do not hesitate to contact the editors with information about your plants, conservation projects, field trips, or noteworthy events. Contributors should review the "Instructions to Authors" printed in the March issue of each year. Advertisers should contact the editors. Views expressed in this publication are those of the authors, not the editorial staff.

All correspondence regarding dues, address changes and missing issues should be sent to the Membership Coordinator at the ICPS. Do not send such correspondence to the editors. Checks for subscriptions should be made to the ICPS in US funds. Dues are \$35 for the first year of membership; renewals are \$30 per year.

ICPS, Inc.
PMB 322
1564-A Fitzgerald Drive
Pinole, CA 94564-2229, USA
icps@carnivorousplants.org

President (Interim)	Richard Myers, richard@carnivorousplants.org
Vice President	Bob Ziemer, bob@carnivorousplants.org
Secretary	Cindy Slezak, cindy@carnivorousplants.org
Treasurer	Richard Myers, richard@carnivorousplants.org
Board Member	Chris Teichreb, chris@carnivorousplants.org
Board Member	Brian Barnes, Conservation Director, brian@carnivorousplants.org
Seed Bank Manager	John Brittnacher, john@carnivorousplants.org (see seed bank ad in this issue)

Editors:

Managing Editor, Stephen Davis, 7010 Via Cordura, San Jose, CA 95139, USA, stephen@carnivorousplants.org

Science Editor, Jan Schlauer, jan@carnivorousplants.org

Science Editor, Fernando Rivadavia, Fernando@carnivorousplants.org

Editor, Bob Ziemer, bob@carnivorousplants.org

Graphic Design: Ken Gumiran, kgumiran@bayareanewsgroup.com

Date of effective publication of the June 2009 issue of Carnivorous Plant Newsletter: 29 June 2009.

The ICPS is the International Cultivar Registration Authority (ICRA) for cultivated carnivorous plants according to The International Code For The Nomenclature of Cultivated Plants. Send relevant correspondence to the ICPS, Inc.

PUBLISHER: ICPS, Inc., Pinole, California. Published quarterly with one volume annually. Graphic Production: Ken Gumiran, 2640 Shadelands Drive, Walnut Creek, CA. Printer: Kandid Litho. Logo and masthead art: Paul Milauskas. Dues are \$35 for the first year of membership; renewals are \$30 per year. © 2009 Carnivorous Plant Newsletter. All rights reserved. ISSN #0190-9215